

## N O T I C E

THIS DOCUMENT HAS BEEN REPRODUCED FROM  
MICROFICHE. ALTHOUGH IT IS RECOGNIZED THAT  
CERTAIN PORTIONS ARE ILLEGIBLE, IT IS BEING RELEASED  
IN THE INTEREST OF MAKING AVAILABLE AS MUCH  
INFORMATION AS POSSIBLE

(NASA-TM-81299) A NUMERICAL SOLUTION FOR  
TWO-DIMENSIONAL FREDHOLM INTEGRAL EQUATIONS  
OF THE SECOND KIND WITH KERNELS OF THE  
LOGARITHMIC POTENTIAL FORM (NASA) 10 p  
HC A02/MF A01

N82-12885

CSCI 12A G3/67 08407  
Unclass

# A Numerical Solution for Two-Dimensional Fredholm Integral Equations of the Second Kind With Kernals of the Logarithmic Potential Form

Ralph E. Gabrielsen and Aynur Ünal

November 1981



**NASA**

National Aeronautics and  
Space Administration

United States Army  
Aviation Research  
and Development  
Command



---

# **A Numerical Solution for Two-Dimensional Fredholm Integral Equations of the Second Kind With Kernals of the Logarithmic Potential Form**

---

Ralph E. Gabrielsen, Aeromechanics Laboratory

AVRADCOM Research and Technology Laboratories

Ames Research Center, Moffett Field, California

Aynur Ünal, NASA-Stanford Joint Institute for Aeronautics and Astronautics  
Stanford, California



National Aeronautics and  
Space Administration

**Ames Research Center**  
Moffett Field, California 94035

United States Army  
Aviation Research and  
Development Command  
St. Louis, Missouri 63166



A NUMERICAL SOLUTION FOR TWO-DIMENSIONAL FREDHOLM  
INTEGRAL EQUATIONS OF THE SECOND KIND WITH KERNELS  
OF THE LOGARITHMIC POTENTIAL FORM

Ralph E. Gabrielsen and Aynur Ünal

Ames Research Center

SUMMARY

Two-dimensional Fredholm integral equations with logarithmic potential kernels are numerically solved. The explicit convergence of these solutions to their true solutions is demonstrated. The results are based on a previous work in which numerical solutions were obtained for Fredholm integral equations of the second kind with continuous kernels.

INTRODUCTION

Previously (ref. 1), the convergence of a numerical scheme for solving one-dimensional Fredholm integral equations of the second kind was proven. Later on (ref. 2), these results were extended to two dimensions for continuous kernels. However, since a class of physical problems involves kernels of the logarithmic potential form, this study extends the theory to kernels of this type.

MAIN DEVELOPMENT

In a recent report (ref. 2), it was shown that the following system of equations,

$$x(t_{ij}) - \lambda \sum_{k=1}^n \sum_{\ell=1}^n h(t_{ij}; t_{k\ell}) x(t_{k\ell}) \Delta = y(t_{ij}) \quad (1)$$

converge to the exact solution of the two-dimensional Fredholm integral equation of the second kind,

$$x(r,s) - \lambda \int_0^1 \int_0^1 h(r,s;t_1,t_2) x(t_1,t_2) dt_1 dt_2 = y(r,s) \quad (2)$$

when the kernel  $h(r,s;t_1,t_2)$  is a continuous function over the unit square  $[0,1] \times [0,1]$  and  $y(r,s)$  is also a continuous function over the unit square.

Equation (2) will be regarded as a functional equation in the Banach space  $X = C^0$  of continuous functions on the unit square  $[0,1] \times [0,1]$  and, typically, will be expressed in the following form:

$$Kx \equiv x - \lambda Hx = y \quad (3)$$

The system (1) is regarded as an approximate functional equation in the space  $\bar{X} = R^n$  and typically is expressed in the following form:

$$\bar{K}\bar{x} \equiv \bar{x} - \lambda \bar{H}\bar{x} = \phi y \quad (4)$$

Let  $\tilde{X}$  be a subspace of  $X$ . Define the mapping  $\phi_0$  in  $\tilde{X}$  onto  $\bar{X}$  as follows: if  $\tilde{x} \in \tilde{X}$ , then  $\phi_0 \tilde{x} = \bar{x}$ .

The differences between this work and that of reference 2 are due to differences arising in proving the following three conditions:

$$\forall \tilde{x} \in \tilde{X}, \quad \|\phi H\tilde{x} - \bar{H}\phi_0 \tilde{x}\| \leq \zeta \|\tilde{x}\| \quad (5)$$

$$\forall x \in X, \quad \exists \tilde{x} \in \tilde{X} \ni \|Hx - \tilde{x}\| \leq \zeta_1 \|x\| \quad (6)$$

$$\exists \tilde{y} \in \tilde{X} \ni \|y - \tilde{y}\| \leq \zeta_2 \|y\| \quad (7)$$

and in showing that  $\zeta$ ,  $\zeta_1$ , and  $\zeta_2$  go to zero as the mesh size goes to zero.

Conditions (5)-(7) are shown for the case of continuous kernels in reference 2. In this work, for the case of logarithmic kernels,

Let  $\bar{H}$  of equation (4) be defined as:

$$\bar{H} \equiv \Delta \begin{bmatrix} 0 & h(t_{11}, t_{12}) \dots h(t_{11}, t_{1n}) \dots h(t_{11}, t_{nn}) \\ h(t_{12}, t_{11}) & 0 & & & h(t_{12}, t_{nn}) \\ \vdots & & & & \vdots \\ h(t_{1n}, t_{11}) & & & \dots & h(t_{1n}, t_{nn}) \\ \vdots & & & & \vdots \\ h(t_{nn}, t_{11}) & h(t_{nn}, t_{12}) \dots h(t_{nn}, t_{nn-1}) & & & 0 \end{bmatrix} \quad (8)$$

*Lemma 1:*  $H$  maps  $C^0 \rightarrow C^0$ .

*Proof of Lemma 1:* Since  $H$  is a bounded linear operator, this result is immediate.

In  $\tilde{X}$  (a subspace of  $X$ ),

$$\tilde{K}\tilde{x} \equiv \tilde{x} - \lambda H\tilde{x} = Py \quad (9)$$

which is the approximate equation as in reference 2, since

$$\phi_0^{-1}\phi = P$$

Equation (9) becomes

$$I - \lambda H\tilde{x} = \phi_0^{-1}\phi y \quad (10)$$

Also,

$$\phi_0(\tilde{x} - \lambda H\tilde{x}) = \phi y \quad (11)$$

$$\bar{x} - \lambda H\bar{y} = \phi y \quad (12)$$

where  $\bar{x} = \phi_0\tilde{x}$ .

*Lemma 2:* Condition (5) is satisfied; that is,

$$\forall \tilde{x} \in \tilde{X}, \quad \|\phi H\tilde{x} - \bar{H}\phi_0\tilde{x}\| \leq \zeta \|\tilde{x}\| \quad (13)$$

*Proof of Lemma 2:* For a given  $\epsilon > 0$ ,  $\exists$  an integral operator  $H_1$  with a continuous kernel  $\mathcal{D}$  (see the appendix),

$$\|H_1 - H\| < \epsilon \quad (14)$$

Since  $H_1$  is an integral operator with a continuous kernel, the results of reference 2 are applicable. In particular,

$$\exists \bar{H}_1 \ni \forall \tilde{x} \in \tilde{X}, \quad \|\phi H_1\tilde{x} - \bar{H}_1\phi_0\tilde{x}\| \leq \zeta_1 \|\tilde{x}\| \quad (15)$$

Hence,

$$\|\phi H\tilde{x} - \bar{H}_1\phi_0\tilde{x}\| \leq \|\phi H\tilde{x} - \phi H_1\tilde{x}\| + \|\bar{H}_1\phi_0\tilde{x} - \bar{H}\phi_0\tilde{x}\| + \|\phi H_1\tilde{x} - \bar{H}_1\phi_0\tilde{x}\| \quad (16)$$

$$\begin{aligned} &\leq \|\phi\| \|\bar{H}_1 - \bar{H}\| \|\tilde{x}\| + \|\bar{H}_1 - \bar{H}\| \|\phi_0\| \|\tilde{x}\| \\ &+ \|\phi H_1\tilde{x} - \bar{H}_1\phi_0\tilde{x}\| \end{aligned}$$

$$\leq \zeta \|\tilde{x}\| \quad (17)$$

Lemma 3: Condition (6) is satisfied; that is,

$$\forall x \in X, \quad \exists \tilde{x} \in \tilde{X} \ni \|Hx - \tilde{x}\| \leq \bar{\zeta}_1 \|x\| \quad (18)$$

Proof of Lemma 3: From the proof of Lemma 2, we know that

$$\forall x \in X, \quad \exists \tilde{x} \in \tilde{X} \ni \|H_1 x - \tilde{x}\| \leq \zeta_1 \|x\| \quad (19)$$

Again, from the proof of Lemma 2,  $\exists H_1 \ni$  for a given  $\epsilon > 0$ ,

$$\|H_1 - H\| < \epsilon \quad (20)$$

Noting that

$$Hx - \tilde{x} = (Hx - H_1 x) + H_1 x - \tilde{x} \quad (21)$$

Now, applying Schwarz's inequality,

$$\|Hx - \tilde{x}\| \leq \|Hx - H_1 x\| + \|H_1 x - \tilde{x}\| \quad (22)$$

$$\leq \|H - H_1\| \|x\| + \|H_1 x - \tilde{x}\| \quad (23)$$

$$\leq \bar{\zeta}_1 \|x\| \quad (24)$$

Condition (7) follows readily from Lemma 3.

From the proofs of Lemmas 2 and 3, it readily follows that  $\zeta$ ,  $\zeta_1$ , and  $\zeta_2$  tend to zero as the mesh size of the partition tends to zero.

#### CONCLUSION

For a given  $\epsilon > 0$ , the solution  $x^*$  of the Fredholm integral equation  $x - \lambda Hx = y$  on  $[0,1] \times [0,1]$  with a logarithmic kernel for continuous  $y$  can be approximated by a function  $\bar{x}^* \ni \|x - \phi_0^{-1} \bar{x}^*\| < \epsilon$ ;  $\bar{x}^*$  satisfies an equation of the form  $\bar{x} - \lambda H\bar{x} = \phi y$ . Hence, the desired accuracy can be achieved by appropriately restricting the mesh size  $\Delta$ .

# APPENDIX

## CONSTRUCTION OF APPROXIMATE INTEGRAL OPERATOR $H_1$

Let  $H$  be an integral operator defined as follows:


$$H[u] = \int_D \log \frac{u}{PQ} [ ]_Q dS_Q$$

*Lemma:* Given  $\epsilon > 0$ ,  $\exists$  an integral operator  $H_1$  with a continuous kernel  $\rho$   $\|H_1 - H\| < \epsilon$ .

*Proof:*

$$H[u] = \int_D \log \frac{r}{PQ} [ ]_Q dS_Q$$

$$D = D_0 \cup D_{r_\epsilon}$$



$$r = |\bar{P} - \bar{Q}|$$

$$= \int_{D_0} \log \frac{r}{PQ} [ ]_Q dS_Q + \int_{D_{r_\epsilon}} \log \frac{r}{PQ} [ ]_Q dS_Q$$

In  $D_0$ , define

$$H_1[u] = \int_{D_0} \log \frac{r}{PQ} [ ]_Q dS_Q$$

In  $D_{r_\epsilon}$ ,

$$H[u] = \int_{D_{r_\epsilon}} \log \frac{r}{PQ} [ ]_Q dS_Q$$



In  $D_{r_\epsilon}$ , define

$$\begin{aligned} H_1[\ ] &= \int_{D_{r_\epsilon}} \log r_\epsilon [\ ] dS \\ &= \log r_\epsilon \int_{D_{r_\epsilon}} [\ ] dS \end{aligned}$$

$$\begin{aligned} \therefore H[\ ] - H_1[\ ] &= \int_{D_{r_\epsilon}} \log r [\ ] dS_Q - \int_{D_{r_\epsilon}} \log r_\epsilon [\ ] dS_Q \\ &= \int_{D_{r_\epsilon}} [\log r - \log r_\epsilon] [\ ] dS_Q \end{aligned}$$

$$\|(H - H_1)[\ ]\| \leq \|[\ ]\| \left( \int_{D_{r_\epsilon}} \log r dS_Q - \int_{D_{r_\epsilon}} \log r_\epsilon dS_Q \right)$$

$$\int_{D_{r_\epsilon}} \log r_\epsilon dS = (\log r_\epsilon) \frac{r_\epsilon^2}{2}$$

$$\begin{aligned} |\log r - \log r_\epsilon| dS &\leq \int \log r_\epsilon dS \\ &\leq \frac{r_\epsilon^2}{2} \log r_\epsilon \end{aligned}$$

$$\therefore \|H - H_1\| \leq \frac{r_\epsilon^2}{2} \log r_\epsilon$$

## REFERENCES

1. Gabrielsen, R. E.: A Solution to the One-Dimensional Fredholm Integral Equations of the Second Kind. NASA TM-81195, 1980.
2. Gabrielsen, R. E.; and Ünal, A.: A Solution for Two-Dimensional Fredholm Integral Equations of the Second Kind with Periodic, Semi-Periodic, and Non-Periodic Kernels. NASA TM-81249, 1981.
3. Kantorovich, V.; and Akilov, G. P.: Functional Analysis in Normed Spaces. Pergamon Press, New York, N. Y., 1964.
4. Kellogg, O. D.: Foundations of Potential Theory. Dover Publications, Inc., New York, 1953.
5. Evans, G. C.: Logarithmic Potential. New York, 1927.